

$$d(x, y) \leq d(x, z) + d(z, y)$$



$$t = d(x, y) - d(y, z) \leq d(x, z)$$

$$-t = d(z, y) - d(y, x) \leq d(x, z)$$


$$|t| = \max\{t, -t\}$$

$$|t| \leq d(x, z)$$



1. Let (M, d) be a metric space.

- (4)  For $x, y, z \in M$ prove that $|d(x, y) - d(y, z)| \leq d(x, z)$.
- (5)  Suppose that $x_n \rightarrow x$ and $y_n \rightarrow y$ in M . Prove that $d(x_n, y_n) \rightarrow d(x, y)$.




2. Let (M, d) be a metric space and let $A, B \subseteq M$.

- (3)  Give the definition of $\text{int}(A)$.

Prove or give a counterexample to the following statements:

- (6)  $\text{int}(A) \cup \text{int}(B) \subseteq \text{int}(A \cup B)$. *mag je $x \in \text{int}(A) \Leftrightarrow \exists r > 0: B_r(x) \subseteq A$ gebruiken*
- (6)  $\text{int}(A \cup B) \subseteq \text{int}(A) \cup \text{int}(B)$.

3. Let (M, d) and (N, ρ) be metric spaces and let $f: M \rightarrow N$ be a function.

- (3)  Complete the following definition: f is *uniformly continuous* if ...
- (6)  Suppose that f is uniformly continuous and let $(x_n)_{n \geq 1}$ be a Cauchy sequence in M . Show that $(f(x_n))_{n \geq 1}$ is a Cauchy sequence in N .
- (7)  Suppose that f is bijective and both f and f^{-1} are uniformly continuous. If N is complete, show that M is complete.

In the next exercise, the following theorem from the book may be useful:



Theorem 8.9 Let (M, d) be a metric space. The following are equivalent:

(i) M is compact.



(ii) If \mathcal{G} is a collection of open sets in M with $M \subseteq \bigcup\{G : G \in \mathcal{G}\}$, then there are finitely many sets $G_1, \dots, G_n \in \mathcal{G}$ such that $M \subseteq \bigcup_{k=1}^n G_k$.

4. Let (M, d) be a metric space. For each $n \geq 1$, let $f_n: M \rightarrow \mathbb{R}$ be a continuous function. Assume that $f_n(x) \rightarrow 0$ for all $x \in M$. Fix $\varepsilon > 0$ and define for $n \geq 1$

$$G_n := \{x \in M : |f_n(x)| < \varepsilon\}.$$

- (4)  a. Prove that G_n is open for all $n \geq 1$. *$\forall n? \quad \varepsilon = r \neq \varepsilon \infty$*
- (4)  b. Prove that $M \subseteq \bigcup_{n=1}^{\infty} G_n$.

Now, in addition, suppose that M is compact and $|f_m(x)| \leq |f_n(x)|$ for all $m \geq n$ and $x \in M$.

- (6)  c. Show that there is an $N \geq 1$ such that $M \subseteq G_N$.
- (4)  d. Prove that $f_n \rightarrow 0$ uniformly on M .

See also the next page.

5. Let (M, d) be a metric space.

(3) Complete the following definition: a set $A \subseteq M$ is *totally bounded* if ...

(9) Let ℓ^1 be the space of all sequences $(x_n)_{n \geq 1}$ such that $\|(x_n)_{n \geq 1}\|_1 := \sum_{n=1}^{\infty} |x_n| < \infty$. Show that

$$A := \{(x_n)_{n \geq 1} : \|(x_n)_{n \geq 1}\|_1 \leq 1\} \subseteq \ell^1$$

$\in / \subseteq ?$
bdd

is bounded, but not totally bounded.

The value of each part of a problem is printed in the margin; the final grade is calculated using

$$\text{Grade} = \frac{\text{Total}}{70} \cdot 9 + 1$$

and rounded in the standard way.

This exam has been composed by the teacher and reviewed by the co-teacher.